



Anti-Human Flt-1 / VEGFR-1, bt (AB-FXG)

Synonyms: vascular endothelial growth factor receptor-1, FLT1, VEGF receptor 1

PLEASE NOTE: ALWAYS CENTRIFUGE VIAL BEFORE OPENING

Size	Order#	Lot#	Expiry Date
50 µg	2033.551.050		
Please enquire for bulk quantities and other vial sizes.			

Description

Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes, dendritic cells and on trophoblast cells. The flt-1 gene was first described in 1990. The receptor contains seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular split tyrosine kinase domain. Compared to VEGFR-2 the Flt-1 receptor has a higher affinity for VEGF but a weaker signaling activity. VEGFR-1 thus leads not to proliferation of endothelial cells, but mediates signals for differentiation. Interestingly a naturally occurring soluble variant of VEGFR-1 (sVEGFR-1) was found in HUVEC supernatants in 1996, which is generated by alternative splicing of the flt-1 mRNA.

- **Source** Mouse
- **Isotype** IgG1
- **Clone** AB-FXG

Biological Activity

ELISA: Use 1–10 µg/ml; Western blotting: Use 2–5 µg/ml; Immunoprecipitation: Use 2-10 µg/ml

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Usage: For research use only. Not for use in diagnostic or therapeutic procedures. Not for human use.

*The Buffer may vary depending on the Lot #. Please contact our technical support if you have specific requirements.

ORDERING
Tel.: +49 40 4320 8448 - 0
order@active-bioscience.de
www.active-bioscience.de

TECHNICAL SUPPORT
Tel.: +49 40 4320 8448 - 11
support@active-bioscience.de

Active Bioscience GmbH
Oberaltenallee 8
D-22081 Hamburg
HRB 98170 Amtsgericht Hamburg