

Murine MANF

Synonyms: Mesencephalic astrocyte-derived neurotrophic factor, Arginine-rich protein, Protein ARMET, Manf, Armet.

PLEASE NOTE: ALWAYS CENTRIFUGE VIAL BEFORE OPENING

Size	Order #	Lot #	Expiry Date
5 µg	3065.960.005		
20 µg	3065.960.020		
1 mg	3065.960.199		

Please enquire for bulk quantities and other vial sizes

Description

MANF Murine Recombinant produced in *E.Coli* is a single, non-glycosylated, polypeptide chain containing 158 amino acids and having a molecular mass of 18.2kDa.

Biological Activity ≥ 100 IU/mg
Source E. Coli

• Purity ≥ 98 % (SDS-PAGE, RP-HPLC)

• Endotoxin level $\leq 0.1 \text{ng/µg} (\leq 1 \text{EU/µg})$

Buffer Lyophilized from a 0.2μm filtered concentrated solution in PBS, pH 7.4*

Physical state
Sterile filtered, lyophilized

Biological Activity

The ED $_{50}$ as determined by a cell proliferation assay using rat C6 cells is less than $10\mu g/ml$, corresponding to a specific activity of >100 IU/mg.

Reconstitution

It is recommended to reconstitute the lyophilized MANF in sterile 18M-cm H2O not less than $100\mu g/ml$, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized MANF although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution MANF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). **Please avoid repeated freeze-thaw cycles**.

Amino Acid Sequence

LRPGDCEVCI SYLGRFYQDL KDRDVTFSPA TIEEELIKFC REARGKENRL CYYIGATDDA ATKIINEVSK PLAHHIPVEK ICEKLKKKDS QICELKYDNQ IDLSTVDLKK LRVKELKKIL DDWGEMCKGC AEKSDYIRKI NELMPKYAPK AASARTDL

Usage: For research use only. Not for use in diagnostic or therapeutic procedures. Not for human use.

^{*}The Buffer may vary depending on the Lot #. Please contact our technical support if you have specific requirements.