

Porcine HGF promoting

Synonyms: Scatter Factor (SF), Hepatopoietin (HPTA), HGF, HGFB, F-TCF, pHGF.

PLEASE NOTE: ALWAYS CENTRIFUGE VIAL BEFORE OPENING

Size	Order #	Lot #	Expiry Date
2 µg	1468.929.002		
10 µg	1468.929.010		
1 mg	1468.929.199		

Please enquire for bulk quantities and other vial sizes

Description

Hepatocyte Growth Factor porcine is extracted from pig liver.

- **Source** Pig Liver
- **Purity** ≥ 98 % (SDS-PAGE, RP-HPLC)
- **Endotoxin level** ≤ 0.1ng/µg (≤ 1EU/µg)
- **Buffer** The sterile protein powder is lyophilized with no additives*
- **Physical state** Sterile filtered, lyophilized

Reconstitution

We recommend a quick spin followed by reconstitution in sterile water to a concentration of at least 100 µg/ml, which can then be further diluted to other aqueous solutions. Do not vortex.

Stability

Lyophilized pHGF although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution pHGF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). **Please avoid repeated freeze-thaw cycles.**

Usage: For research use only. Not for use in diagnostic or therapeutic procedures. Not for human use.

*The Buffer may vary depending on the Lot #. Please contact our technical support if you have specific requirements.

ORDERING
Tel.: +49 40 43208448-0
order@active-bioscience.de
www.active-bioscience.de

TECHNICAL SUPPORT
Tel.: +49 40 43208448-11
support@active-bioscience.de

Active Bioscience GmbH
Oberaltenallee 8
D-22081 Hamburg
HRB 98170 Amtsgericht Hamburg